



ACC.14

TCT@ACC-12 | innovation in intervention

A27

JACC April 1, 2014

Volume 63, Issue 12



Acute Coronary Syndromes

PREDICTORS ASSOCIATED WITH ACUTE STENT THROMBOSIS AFTER PRIMARY PCI: THE EUROMAX TRIAL

Oral Contributions

Room 147 B

Saturday, March 29, 2014, 9:15 a.m.-9:30 a.m.

Session Title: Antiplatelet Therapy in ACS: Challenges and Controversies

Abstract Category: 3. Acute Coronary Syndromes: Therapy

Presentation Number: 901-08

Authors: *Peter Clemmensen, Arnoud van't Hof, Efthymios N. Deliargyris, Pierre Coste, Jurrien ten Berg, Claudio Cavallini, Martial Hamon, Dariusz Dudek, Uwe Zeymer, Xavier Tabone, Tim Clayton, Debra Bernstein, Jayne Prats, Philippe Gabriel Steg, Rigshospitalet, Copenhagen, Denmark, Université Paris-Diderot and Assistance Publique-Hôpitaux de Paris, Paris, France*

Background: Acute stent thrombosis (AST) after primary PCI (pPCI) for ST-segment elevation myocardial infarction (STEMI) is a rare event occurring in $\leq 1\%$ of patients. We determined clinical, procedural and treatment related factors associated with AST in EUROMAX.

Methods: EUROMAX included 2198 patients transported for pPCI. Baseline, clinical and procedural characteristics and treatment strategies [P2Y₁₂ inhibitor choice, bivalirudin with reduced dose post-PCI infusion (0.25 mg/kg/h, BIV-Lo) vs. bivalirudin with prolonged PCI dose (1.75 mg/kg/h, BIV-PCI) vs. heparins \pm GPI] were compared between patients with or without AST using the Chi-Square test. Logistic regression was performed using a forward stepwise procedure to select candidate variables.

Results: The overall rate of AST was 0.6% and was higher with bivalirudin vs. heparins \pm GPI (1.1% vs. 0.2%; $P = 0.007$). The median time for AST was 2.3 hours (IQR 1.9-2.8). Patients with AST vs. without AST had less hypertension [2/14, (14%); 961/2182 (44.0%), $p=0.03$], and were more likely to have received GPI [11/14 (78.6%); 880/2183 (40.3%), $p=0.004$]. Of the patients in the bivalirudin arm with AST, 3 did not receive a prolonged infusion and among the 9 who did, 8 received BIV-Lo and 1 BIV-PCI. Multivariate analysis found hypertension (OR 0.21, $p=0.047$) and receipt of BIV-Lo (OR 7.7, $p=0.009$) as predictors of AST. Incidence of AST in BIV-PCI as compared with heparins \pm GPI was [1/191 (0.5%) vs 2/947 (0.2%), $p=0.46$]. The rate of AST in BIV-Lo was significantly higher than heparins \pm GPI [8/659 (1.2%) vs 2/947 (0.2%), $p=0.03$]. Oral P2Y₁₂ inhibitor choice (clopidogrel vs. prasugrel vs. ticagrelor) had no impact on AST.

Conclusion: This post-hoc analysis from EUROMAX suggests that AST happens in the first few hours after pPCI and that neither the new oral P2Y₁₂ inhibitors (prasugrel or ticagrelor) nor a low-dose (0.25 mg/kg/h) bivalirudin infusion are protective. However, prolonging bivalirudin infusion at the PCI dose was not associated with a higher risk of AST.